











unsealed controls. Annualized caries progression over 3 years found to be reduced by 73.2% among sealed tooth compared to unseal. GC Fuji VII may be a better choice for preventing caries in children from low socioeconomic, high caries risk population in rural set up to have a caries-free generation.

### Acknowledgement

We express our whole hearted thanks to JSS Academy of Higher Education and Research, Mysuru for funding this project and JSS Free Residential School, Suttur, authorities for giving permission for carrying out this study.

### Financial support and sponsorship

This study was financially supported by university grants from JSS Academy of Higher Education and Research, Mysuru, India.

### Conflicts of interest

There are no conflicts of interest.

### REFERENCES

- Locker D. Deprivation and oral health: A review. *Community Dent Oral Epidemiol* 2000;28:161-9.
- Watt RG, Listl S, Peres M, Heilmann A. Social inequalities in oral health: From evidence to action. *Int Centre Oral Health Inequal Res Policy* 2015;1-44. Available from: [http://media.news.health.ufl.edu/misc/cod-ralhealth/docs/posts\\_frontpage/SocialInequalities.pdf](http://media.news.health.ufl.edu/misc/cod-ralhealth/docs/posts_frontpage/SocialInequalities.pdf). [Last accessed on 2020 May 29].
- Splieth C, Förster M, Meyer G. Additional caries protection by sealing permanent first molars compared to fluoride varnish applications in children with low caries prevalence: 2-year results. *Eur J Paediatr Dent* 2001;2:133-8.
- Damle SC, Patel AR. Caries prevalence and treatment need amongst children of Dharavi, Bombay, India. *Community Dent Oral Epidemiol* 1994;22:62-3.
- Tewari A, Chawla HS. Study of prevalence of dental caries in an urban area of India (Chandigarh). *J Indian Dent Assoc* 1977;49:231-9.
- Dhar V, Jain A, Van Dyke TE, Kohli A. Prevalence of dental caries and treatment needs in the school-going children of rural areas in Udaipur district. *J Indian Soc Pedod Prev Dent* 2007;25:119-21.
- Saravanan S, Kalyani V, Vijayarani MP, Jayakodi P, Felix J, Arunmozhi P, *et al.* Caries prevalence and treatment needs of rural school children in Chidambaram Taluk, Tamil Nadu, South India. *Indian J Dent Res* 2008;19:186-90.
- Bali RK, Mathur VB, Talwar PP, Channa HB. National Oral Health Survey & Fluoride Mapping. New Delhi: Dental Council of India; 2002.
- McDonald SP, Sheiham A. The distribution of caries on different tooth surfaces at varying levels of caries – A compilation of data from 18 previous studies. *Community Dent Health* 1992;9:39-48.
- Brown LJ, Selwitz RH. The impact of recent changes in the epidemiology of dental caries on guidelines for the use of dental sealants. *J Public Health Dent* 1995;55:274-91.
- Zenkner JE, Alves LS, de Oliveira RS, Bica RH, Wagner MB, Maltz M. Influence of eruption stage and biofilm accumulation on occlusal caries in permanent molars: A generalized estimating equations logistic approach. *Caries Res* 2013;47:177-82.
- Alves LS, Zenkner JE, Wagner MB, Damé-Teixeira N, Susin C, Maltz M. Eruption stage of permanent molars and occlusal caries activity/arrest. *J Dent Res* 2014;93:114S-9.
- Dye BA, Thornton-Evans G, Li X, Iafolla TJ. Dental caries and sealant prevalence in children and adolescents in the United States, 2011–2012. *NCHS Data Br* 2015;191:1-8.
- Wright JT, Crall JJ, Fontana M, Gillette EJ, Nový BB, Dhar V, *et al.* Evidence-based clinical practice guideline for the use of pit-and-fissure sealants: A report of the American Dental Association and the American Academy of Pediatric Dentistry. *J Am Dent Assoc* 2016;147:672-82.
- Gooch BF, Griffin SO, Gray SK, Kohn WG, Rozier RG, Siegal M, *et al.* Preventing dental caries through school-based sealant programs, updated recommendations and reviews of evidence. *J Am Dent Assoc* 2009;140:1356-65.
- Truman BI, Smith-Akin CK, Hinman AR, Gebbie KM, Brownson R, Novick LF, *et al.* Developing the guide to community preventive services – Overview and rationale. The task force on community preventive services. *Am J Prev Med* 2000;18:18-26.
- Yang S. Evaluation of the effect of pit and fissure sealing. *J Dent Dis Prev* 1994;2:41-2.
- Tonn EM, Ryge G. Three year clinical evaluation of four sealants in Los Altos, CA. *J Dent Res* 1982;61:331.
- Sly EG, Kalpana AE, Missana L. Clinical evaluation of glass ionomer for pit and fissure sealing of fully erupted molars. *Acta Odontol Latinoam* 2010;23:3-7.
- Ninawe N, Ullal NA, Khandelwal V. A 1-year clinical evaluation of fissure sealants on permanent first molars. *Contemp Clin Dent* 2012;3:54-9.
- Guler C, Yilmaz Y. A two-year clinical evaluation of glass ionomer and ormocer based fissure sealants. *J Clin Pediatr Dent* 2013;37:263-8.
- Chen XX, Liu XG. Clinical comparison of Fuji VII and a resin sealant in children at high and low risk of caries. *Dent Mater J* 2013;32:512-8.
- Haznedarog̃lu E, Güner S, Duman C, Menteş A. A 48-month randomized controlled trial of caries prevention effect of a one-time application of glass ionomer sealant versus resin sealant. *Dent Mater J* 2016;35:532-8.
- Al-Jobair A, Al-Hammad N, Alsadhan S, Salama F. Retention and caries-preventive effect of glass ionomer and resin-based sealants: An 18-month-randomized clinical trial. *Dent Mater J* 2017;36:654-61.
- Prathibha B, Reddy PP, Anjum MS, Monica M, Praveen BH. Sealants revisited: An efficacy battle between the two major types of sealants – A randomized controlled clinical trial. *Dent Res J (Isfahan)* 2019;16:36-41.
- Ganesh M, Tandon S. Clinical evaluation of FUJI VII sealant material. *J Clin Pediatr Dent* 2006;31:52-7.
- Kamala BK, Hegde AM. Fuji III vs. Fuji VII glass ionomer sealants – A clinical study. *J Clin Pediatr Dent* 2008;33:29-33.
- Yazici AR, Kiremitçi A, Celik C, Ozgünaltay G, Dayangaç B. A two-year clinical evaluation of pit and fissure sealants placed with and without air abrasion pretreatment in teenagers. *J Am Dent Assoc* 2006;137:1401-5.
- Liu W, Xiong L, Li J, Guo C, Fan W, Huang S. The anticaries effects of pit and fissure sealant in the first permanent molars of school-age children from Guangzhou: A population-based cohort study. *BMC Oral Health* 2019;19:156.
- Williams B, Laxton L, Holt RD, Winter GB. Fissure sealants: A 4-year clinical trial comparing an experimental glass polyalkenoate cement with a bis glycidyl methacrylate resin used as fissure sealants. *Br Dent J* 1996;180:104-8.
- Trairatvorakul C, Kladaew S, Songsiripradaboon S. Active management of incipient caries and choice of materials. *J Dent Res* 2008;87:228-32.
- Bayrak S, Tunc ES, Aksoy A, Ertas E, Guvenc D, Ozer S. Fluoride release and recharge from different materials used as fissure sealants. *Eur J Dent* 2010;4:245-50.
- Gokhale N, Nuvvula S. Influence of socioeconomic and working status of the parents on the incidence of their children's dental caries. *J Nat Sci Biol Med* 2016;7:127-9.
- Touger-Decker R, Mobley C. Academy of Nutrition and Dietetics. Position of the Academy of Nutrition and Dietetics: Oral health and nutrition. *J Acad Nutr Diet* 2013;113:693-701.