

Dens invagination: A review of literature and report of two cases

**Seema Thakur,
Narbir S. Thakur¹,
Manmohan Bramta²,
Mohit Gupta³**

Departments of Pediatric and Preventive Dentistry, ¹Oral Pathology, ²Oral and Maxillofacial Surgery, H.P. Government Dental College, Shimla, Himachal Pradesh, ³Private Practitioner Orthodontist, Jammu, India

Address for correspondence:

Dr. Seema Thakur, Department of Paediatric and Preventive Dentistry, H.P. Government Dental College, Shimla, Himachal Pradesh, India. E-mail: cima2009@hotmail.com

Abstract

Dens invaginatus occurs as a result of the invagination of the enamel organ. These cases may present difficulties with respect to its diagnosis and treatment because of canal morphology. It frequently leads to caries, pulpal, and periodontal involvement with necrosis and loss of attachment. The knowledge of classification and anatomical variations of teeth with dens invaginatus are of great importance for correct treatment. This article presents two case reports of two different types of dens invaginatus along with profound review of the literature regarding etiology, epidemiology, and histology. It discusses clinical appearance and diagnosis, and it provides guidelines for decision-making and treatment of invaginated teeth.

Key words: Case reports, classification, dens invaginatus, Etiology

INTRODUCTION

Dens invaginatus is a developmental anomaly resulting from the invaginations of the enamel organ into the dental papilla during the soft tissue stage of development. As the hard tissues are formed, the invaginated enamel organ produces a small tooth within the future pulp chamber. Dens invagination in a human tooth was first described by a dentist named ‘Socrates’ in 1856.^[1]

Etiology

The etiology of dens invaginatus malformation is controversial and remains unclear. Over the last decades, several theories have been proposed to explain the etiology of dental coronal invaginations

Kronfeld (1934) suggested that the invagination results from a focal failure of growth of the internal enamel epithelium while the surrounding normal epithelium continues to proliferate and engulf the static area.^[2]

Rushton (1937) proposed that the invagination is a result of rapid and aggressive proliferation of a part of the internal enamel epithelium invading the dental papilla.^[3]

Oehlers (1957) considered that distortion of the enamel organ during tooth development and subsequent protrusion of a part of the enamel organ will lead to the formation of an enamel-lined channel ending at the

cingulum or occasionally at the incisal tip. The latter might be associated with irregular crown form.^[4,5]

Atkinson (1943) suggested that the problem was the result of external forces exerting an effect on the tooth germ during development.^[6]

Genetic factor has been proposed to be the cause.^[7,8]

Classification

The first documented attempt to classify dens invaginatus was by Hallett^[9] (1953) who suggested the existence of 4 types of invagination based on both clinical and radiographic criteria. Other classifications have also been described involving a variety of criteria and standards.^[10,11] For example, Schulze and Brand^[12] (1972) suggested an assessment based on 12 possible variations in clinical and radiographic appearance of the invagination. However, the system described by Oehlers^[4] (1957a) appears to be the most widely used, possibly because of its simple nomenclature and ease of application. This system categorizes invaginations into 3 classes as determined by how far they extend radiographically from the crown into the root [Figure 1].

Type I: The invagination is minimal and enamel-lined; it is confined within the crown of the tooth and does not extend beyond the level of the external amelo-cemental junction.

Type II: The invagination is enamel-lined and extends into

the pulp chamber, but remains within the root canal with no communication with the periodontal ligament.

Type III A: The invagination extends through the root and communicates laterally with the periodontal ligament space through a pseudo-foramen. There is usually no communication with the pulp, which lies compressed within the root.

Type III B: The invagination extends through the root and communicates with the periodontal ligament at the apical foramen. There is usually no communication with the pulp.

In Type III lesions, any infection within the invagination can lead to an inflammatory response within the periodontal tissues giving rise to a 'peri-invagination periodontitis.'

The limitations associated with the use of conventional radiography in the classification and management of dens invaginatus may be overcome in the future with the increasing availability of computerized 3D imaging.^[13,14] Currently, such clinical techniques do not provide images of sufficient quality to fully evaluate the morphology of an invagination *in situ* although for extracted teeth sufficient detail can be obtained.

Prevalence and distribution

The reported prevalence of adult teeth affected with dens invaginatus is between 0.3% and 10% with the problem observed in 0.25% to 26.1% of individuals examined [Table 1].

CASE REPORTS

Case report 1

A 12-year-old boy reported with a chief complaint of pain with respect to upper anterior tooth since 3 days. The patient was in good general health. Extra-oral examination revealed no significant findings. Intra-oral examination revealed a comparatively wider maxillary right central incisor with a deep anatomic pit on palatal surface [Figure 2]. The tooth was tender on percussion and did not respond to thermal and electric stimuli. The periapical radiograph revealed type II DI with open apex




Figure 1: Oehler's classification of dens invaginatus (Coronal types)

and periapical pathology with respect to maxillary right central incisor [Figure 3]. Endodontic treatment was started immediately.

Case report 2

A 10-year-old boy reported with a chief complaint of non-eruption upper right anterior tooth. The patient was in good general health. Extra-oral examination revealed no significant findings. Intra-oral examination showed fibrotic gingival eruption bulge of maxillary left lateral incisor. The periapical radiograph showed type III DI with respect to unerupted maxillary left lateral incisor [Figure 4]. Keeping in view the age of the patient and that the contralateral maxillary right lateral incisor was erupted well, it was decided that an incision should be made on the eruption bulge. The incision was made, and patient was reviewed after 1 month. After eruption, the prophylactic composite sealant was applied on the invagination.

DISCUSSION

In most cases, a dens invaginatus is detected by chance on the radiograph. Clinically, unusual crown morphology ('dilated,' 'peg-shaped,' 'barrel-shaped') or a deep foramen coecum may be important hints, but affected teeth also may show no clinical signs of the malformation. As pulpal involvement of teeth with coronal invaginations may occur a short time after tooth eruption, an early diagnosis is mandatory to instigate preventive treatment.^[31]

In the first case report, there was unusual crown morphology of the tooth with pulpal involvement, whereas the second case was detected radiographically in the unerupted tooth.

The invagination allows entry of irritants into an area,



Figure 2: Palatal surface of maxillary right central incisor of first case

Table 1: Prevalence studies on dens invaginatus

Authors	Year	Sample	Frequency
Muhlreiter ^[15]	1873	500 maxillary lateral incisors	2.8%
Atkinson ^[6]	1943	500 maxillary lateral incisors	10% of teeth
Boyne ^[16]	1952	1000 maxillary incisors	8%
Shafer ^[17]	1953	2542 Full-mouth surveys	1.3% of patients
Hallet ^[9]	1953	586 Full-mouth surveys	6.6% of lateral incisor 0.5% of central incisors
Amos ^[18]	1955	1000 Full-mouth surveys	5.1% of patients
Amos ^[18]	1955	203 Full-mouth surveys	6.9% of students of dentistry
Hermel ^[10]	1964s	500 Full-mouth surveys	2% of patients
Poyton and Morgan ^[19]	1966	5000 Full-mouth surveys	0.25% of patient
Miyoshi <i>et al.</i> ^[20]	1971	Extracted maxillary lateral incisors	38.5% of teeth
Fujiki <i>et al.</i> ^[21]	1974	2126 Lateral maxillary incisors	4.2% of teeth
Thomas ^[22]	1974	1886 Full-mouth survey	7.74% of patients
Gotoh <i>et al.</i> ^[23]	1979	766 Maxillary lateral incisors	9.66% of teeth
Ruprecht <i>et al.</i> ^[24]	1986	1581 Full-mouth surveys	1.7% of patients
Ruprecht <i>et al.</i> ^[25]	1987	300 Full-mouth surveys	10% of patients
Thongudomporn and Freer ^[26]	1998	111 Full-mouth surveys	26.1% of patients
Backman and Wahlin ^[27]	2001	739 Full-mouth surveys	6.8% of patients
Hamasha and Al-Omar ^[28]	2004	1660 Full-mouth survey	2.95% of patients and 0.65% of teeth
Ezoddini <i>et al.</i> ^[29]	2007	480 Dental panoramic tomograph	0.8%
Cakici <i>et al.</i> ^[30]	2010	1012 Full-mouth surveys	1.3%

**Figure 3:** Periapical view of dens invaginatus Type II in maxillary right central incisor

which is separated from pulpal tissue by only a thin layer of enamel and dentine and present a predisposition for the development of dental caries. In some cases, the enamel-lining is incomplete. Channels may also exist between the invagination and the pulp.^[3,32] Therefore, pulp necrosis often occurs rather early, within a few years of eruption,^[19,33,34] sometimes even before root end closure. Other reported sequelae of undiagnosed and untreated coronal invaginations are abscess formation,^[35,36] retention of neighboring teeth,^[37,38] cysts,^[35] and internal resorption.^[39]

The treatments options are; prophylactic or preventive sealing of the invagination,^[40,41] root canal treatment,^[42] endodontic apical surgery,^[43,44] intentional replantation,^[45] and extraction.^[46]

**Figure 4:** Periapical view of dens invaginatus Type III in maxillary left lateral incisor

In the first case, endodontic management is required, which includes apexification followed by obturation of the tooth, whereas in the second case, prophylactic sealing of the invagination was done after the eruption of the tooth.

CONCLUSION

The knowledge of classification and anatomic variations of teeth with dens invaginatus is important for early detection and early management for the practitioners.

REFERENCES

- Shulze C. Developmental abnormalities of the teeth and the jaws. In: Gorlin O, Goldman H, editors. Thoma's Oral Pathology. St.Louis;

- Mosby: 1970. p. 96-183.
2. Kronfeld R. Dens in dente. *J Dent Res* 1934;14:49-66.
 3. Rushton MA. A collection of dilated composite odontomas. *Br Dent J* 1937;63:65-85.
 4. Oehlers FA. Dens invaginatus I. Variations of the invagination process and associated anterior crown forms. *Oral Surg Oral Med Oral Pathol* 1957;10:1204-18.
 5. Oehlers FA. Dens invaginatus II. Associated posterior crown forms and pathogenesis. *Oral Surg Oral Med Oral Pathol* 1957;10:1302-16.
 6. Atkinson SR. The permanent maxillary lateral incisor. *Am J Orthod* 1943;29:685-98.
 7. Hosey MT, Bedi R. Multiple dens invaginatus in two brothers. *Endod Dent Traumatol* 1996;12:44-7.
 8. Pokala P, Acs G. A constellation of dental anomalies in a chromosomal deletion syndrome (7q32): Case report. *Pediatr Dent* 1994;16:306-9.
 9. Hallet GE. The incidence, nature and clinical significance of palatal invagination in the maxillary incisors teeth. *Proc R Soc Med* 1953;46:491-9.
 10. Ulmanky M, Hermel J. Double dens in dente in a single tooth. *Oral Surg Oral Med Oral Pathol* 1964;17:92-7.
 11. Vincent-Townend J. Dens invaginatus. *J Dent* 1974;2:234-8.
 12. Schulze C, Brand E. Uber den dens invaginatus (Dens in dente). *ZWR* 1972;81:569-73.
 13. Vannier MW, Hildebolt CF, Conover G, Knapp RH, Yokoyama-Crothers N, Wang G. Three dimensional dental imaging by spiral CT. A progress report. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1997;84:561-70.
 14. Sponchiado EC Jr, Ismail HA, Braga MR, de Carvalho FK, Simoes CA. Maxillary central incisor with two root canals: A case report. *J Endod* 2006;32:1002-4.
 15. Muhleirer E. Die natur der anomalen hohlenbildung imoberen Seitenschneidezahne. *Deutsche vierteljahresschrift für zahnkunde* 1873;13:367-72.
 16. Boyne PJ. Dens in dente: Report of three cases. *J Am Dent Assoc* 1952;45:208-9.
 17. Shafer WG. Dens in dente. *NY State Dent J* 1953;19:220-5.
 18. Amos ER. Incidence of the small dens in dente. *J Am Dent Assoc* 1955;51:31-3.
 19. Poyton HG, Morgan GA. Dens in dente. *Dent Radiogr Photogr* 1966;39:27-33.
 20. Miyoshi S, Fujiwara J, Nakata T, Yamamoto K, Deguchi K. Dens invaginatus in Japanese incisors. *Shika Kiso Igakkai Zasshi* 1971;13:539-44.
 21. Fujiki Y, Tamaki N, Kawahara K, Nabae M. Clinical and radiographic observations of dens invaginatus. *Dentomaxillofac Radiol* 1974;3:343-8.
 22. Thomas JG. A study of dens in dente. *Oral Surg Oral Med Oral Pathol* 1974;38:653-5.
 23. Gotoh T, Kawahara K, Imai K, Kishi K, Fujiki Y. Clinical and radiographic study of dens invaginatus. *Oral Surg Oral Med Oral Pathol* 1979;48:88-91.
 24. Ruprecht A, Batniji S, Sastry KA, El-Neweihi E. The incidence of dental invagination. *J Pedod* 1986;10:265-72.
 25. Ruprecht A, Sastry KA, Batniji S, Lambourne A. The clinical significance of dental invagination. *J Pedod* 1987;11:176-81.
 26. Thongudomporn U, Freer TJ. Prevalence of dental anomalies in orthodontic patients. *Aust Dent J* 1998;43:395-8.
 27. Backman B, Wahlin YB. Variations in number and morphology of permanent teeth in 7-year-old Swedish children. *Int J Paediatr Dent* 2001;11:11-7.
 28. Hamasha AA, Al-Omari QD. Prevalence of dens invaginatus in Jordanian adults. *Int Endod J* 2004;37:307-10.
 29. Ezoddini AF, Sheikhha MH, Ahmadi H. Prevalence of dental developmental anomalies: A radiographic study. *Community Dent Health* 2007;24:140-4.
 30. Cakici F, Celiekoglu M, Arslan H, Topcuoglu HS, Erdogan AS. Assessment of the prevalence and characteristics of dens invaginatus in a sample of Turkish Anatolian population. *Med Oral Patol Oral Cir Bucal* 2010;15:e855-8.
 31. Hu Ismann M. Dens invaginatus: Aetiology, classification, prevalence, diagnosis, and treatment considerations. *Int Endod J* 1997;30:79-90.
 32. Stamfelj I, Kansky AA, Gaspersic D. Unusual variant of type 3 dens invaginatus in a maxillary canine: A rare case report. *J Endod* 2007;33:64-8.
 33. Hitchin AD, MCHugh WD. Three coronal invaginations in a dilated composite odontome. *Br Dent J* 1954;97:90-2.
 34. Greenfeld RS, Cambuzzi JV. Complexities of endodontic treatment of maxillary lateral incisors with anomalous root formation. *Oral Surg Oral Med Oral Pathol* 1986;62:82-8.
 35. Nik-Hussein NN. Dens invaginatus: Complications and treatment of non-vital infected tooth. *J Clin Pediatr Dent* 1994;18:303-6.
 36. Chen RJ, Yang JF, Chao TC. Invaginated tooth associated with periodontal abscess. *Oral Surg Oral Med Oral Pathol* 1990;69:659.
 37. Whyman RA, Macfadyen EE. Dens in dente associated with infective endocarditis. *Oral Surg Oral Med Oral Pathol* 1974;78:47-50.
 38. Conklin WW. Double bilateral dens invaginatus in the maxillary incisor region. *Oral Surg Oral Med Oral Pathol* 1975;39:949-52.
 39. Mader C. Triple dens in dente. *Oral Surg Oral Med Oral Pathol* 1977;44:966.
 40. Shapiro L. Dens in dente. *Oral Surg Oral Med Oral Pathol* 1970;30:782.
 41. Aguiar CM, Ferreira JP, Câmara AC, de Figueiredo JA. Type 2 dens invaginatus in a maxillary lateral incisor: A case report of a conventional endodontic treatment. *J Clin Pediatric Dent* 2008;33:103-6.
 42. Steffen H, Splieth C. Conventional treatment of dens invaginatus in maxillary lateral incisor with sinus tract: One year follow-up. *J Endod* 2005;31:130-3.
 43. Ortiz P, Weisleder R, Villareal de Justus Y. Combined therapy in the treatment of dens invaginatus: Case report. *J Endod* 2004;30:672-4.
 44. Nallapati S. Clinical management of a maxillary Lateral incisor with vital pulp and type III dens invaginatus; a case report. *J Endod* 2004;30:726-31.
 45. Nedley MP, Power GK. Intentional extraction and reimplantation of an immature invaginated central incisor. *ASDC J Dent Child* 1997;64:417-20.
 46. Girsch WJ, Mc Clammy TV. Microscopic removal of dens invaginatus. *J Endod* 2002;28:336-9.

How to cite this article: Thakur S, Thakur NS, Bramta M, Gupta M. Dens invagination: A review of literature and report of two cases. *J Nat Sc Biol Med* 2014;5:218-21.

Source of Support: Nil. **Conflict of Interest:** None declared.

Access this article online

Quick Response Code:	Website: www.jnsbm.org
	DOI: 10.4103/0976-9668.127341