

vigilant postoperative monitoring. Keeping this in view, we had monitored the patient closely for 48 h in the Intensive Care Unit.

Epidural anesthesia provides a safer alternative to general anesthesia in parturient with complex valvular lesions. However, it is easier said than done. Successful management necessitates strict vigilance and an extremely cautious approach to maintain the hemodynamic stability throughout the peripartum period.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Datta S, Kodali BS, Segal S. *Obstetric Anaesthesia Handbook*. 5th ed. Boston: Springer; 2010.
- Harnett M, Tsen LC. Cardiovascular disease. In: Chestnut DH, Polley LS, Tsen LC, editors. *Chestnut's Obstetric Anaesthesia: Principles and Practice*. 4th ed. Philadelphia (PA): Mosby Elsevier; 2009. p. 883-4.
- Langesaeter E, Dragsund M, Rosseland LA. Regional anaesthesia for a caesarean section in women with cardiac disease: A prospective study. *Acta Anaesthesiol Scand* 2010;54:46-54.
- Sharma SK, Gambling DR, Gajraj NM, Truong C, Sidawi EJ. Anesthetic management of a parturient with mixed mitral valve disease and uncontrolled atrial fibrillation. *Int J Obstet Anesth* 1994;3:157-62.
- Gash AK, Carabello BA, Kent RL, Frazier JA, Spann JF. Left ventricular performance in patients with coexistent mitral stenosis and aortic insufficiency. *J Am Coll Cardiol* 1984;3:703-11.
- Celik M, Dostbil A, Alici HA, Sevimli S, Aksoy A, Erdem AF, *et al*. Anaesthetic management for caesarean section surgery in two pregnant women with severe pulmonary hypertension due to mitral valve stenosis. *Balkan Med J* 2013;30:439-41.
- Harsoor SS, Joshi SD. Anaesthetic management of parturient with primary pulmonary hypertension posted for caesarean section — A case report. *Indian J Anaesth* 2005;49:223-5.
- Kannan M, Vijayanand G. Mitral stenosis and pregnancy: Current concepts in anaesthetic practice. *Indian J Anaesth* 2010;54:439-44.
- Coskun D, Mahli A, Korkmaz S, Demir FS, Inan GK, Erer D, *et al*. Anaesthesia for caesarean section in the presence of multivalvular heart disease and severe pulmonary hypertension: A case report. *Cases J* 2009;2:9383.
- Weeks SK, Smith JB. Obstetric anaesthesia in patients with primary pulmonary hypertension. *Can J Anaesth* 1991;38:814-6.
- Gomar C, Errando CL. Neuroaxial anaesthesia in obstetrical patients with cardiac disease. *Curr Opin Anaesthesiol* 2005;18:507-12.
- Dyer RA, Butwick AJ, Carvalho B. Oxytocin for labour and caesarean delivery: Implications for the anaesthesiologist. *Curr Opin Anaesthesiol* 2011;24:255-61.
- Secher NJ, Arnsbo P, Wallin L. Haemodynamic effects of oxytocin (syntocinon) and methyl ergometrine (methergin) on the systemic and pulmonary circulations of pregnant anaesthetized women. *Acta Obstet Gynecol Scand* 1978;57:97-103.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Access this article online	
Quick Response Code:	Website: www.jnsbm.org
	DOI: 10.4103/0976-9668.184708

How to cite this article: Sen S, Chatterjee S, Mazumder P, Mukherji S. Epidural anesthesia: A safe option for cesarean section in parturient with severe pulmonary hypertension. *J Nat Sc Biol Med* 2016;7:182-5.